

BLACK HOLE MASS, VELOCITY DISPERSION AND THE RADIO SOURCE IN AGN

CHARLES H. NELSON

Physics Dept., University of Nevada, Las Vegas, Box 4002, 4505 Maryland Pkwy., Las Vegas, NV 89154,
 cnelson@physics.unlv.edu

Draft version February 1, 2008

ABSTRACT

The recent discovery of a correlation between nuclear black hole mass, $M_\bullet$, and the stellar velocity dispersion, σ_* (Gebhardt *et al.* 2000a; Ferrarese & Merritt 2000), in elliptical galaxies and spiral bulges, has raised the question whether such a relationship exists for AGN. Estimates of $M_\bullet$ for many AGN, made using reverberation mapping techniques, allow exploration of the relationship between black hole mass, the host galaxy and the energetics of nuclear emission. However, since only a few AGN have both $M_\bullet$ and σ_* measurements, we use the [OIII] λ 5007 emission line widths on the assumption that for most AGN the forbidden line kinematics are dominated by virial motion in the host galaxy bulge. We find that a relation does exist between $M_\bullet$ and [OIII] line width for AGN which is similar to the one found by Gebhardt *et al.* (2000a), although with more scatter as expected if secondary influences on the gas kinematics are also present. Our conclusion is that both active and inactive galaxies follow the same relationship between black hole mass and bulge gravitational potential. We find no compelling evidence for systematic differences in the mass estimates from reverberation mapping and stellar dynamics. We also find that for radio quiet AGN the radio power and black hole mass are highly correlated linking emission on scales of kiloparsecs with the nuclear energy source.

Subject headings: galaxies: nuclei — galaxies: Seyfert — quasars: general — galaxies: kinematics and dynamics

1. INTRODUCTION

In the standard model for an active galactic nucleus (AGN), radiant energy is released through accretion onto a supermassive ($10^6 - 10^9 M_\odot$) black hole. However, until recently, there have been few mass estimates for nuclear black holes in AGN. Kinematic studies of nuclear gas disks (*e.g.* Harms *et al.* 1994) and stellar dynamical studies, using high spatial resolution ground-based spectroscopy (*e.g.* Kormendy 1988) and more recently with the Space Telescope Imaging Spectrograph (STIS) (*e.g.* Bower *et al.* 2000), have produced numerous central mass estimates mostly for normal galaxies. However, in application to AGN these techniques are severely limited. Since roughly one galaxy in a hundred has a luminous active nucleus even the nearest examples are relatively distant reducing the effective spatial resolution. Furthermore, stellar dynamical techniques require high signal-to-noise measurements of stellar absorption features which are often lost in the glare of a bright active nucleus. Also emission lines from ionized gas at small radii can be influenced by relativistic jets or winds associated with the AGN.

These difficulties can be overcome, however, using reverberation mapping techniques (*e.g.* Netzer & Peterson 1997). Monitoring the variability of the continuum and emission line fluxes in broad line AGN, shows that the two are highly correlated but that the emission lines lag behind the continuum by times ranging from days to months. The lag is best explained as a light travel time effect and can be used to estimate the size of the Broad Line Region (BLR). If one further assumes that the BLR kinematics are largely Keplerian, one can use the line widths as a characteristic velocity and in turn determine the total mass contained within the BLR. Strong evidence for Keplerian motion ex-

ists for at least one well-studied Seyfert, NGC 5548. Results for several different emission lines, each with different lags and therefore spanning a range of distances from the nucleus, give consistent values for the central mass (Peterson & Wandel 1999). Reverberation studies have now yielded $M_\bullet$ estimates for numerous Seyfert 1 galaxies and quasars (*e.g.* Kaspi *et al.* 2000). However, there have been some claims that the masses determined from these variability studies are systematically low (Wandel *et al.* 1999; Ho 1999).

Recent work has focused on the distributions of black hole properties and the relationship to their host galaxies. Kormendy & Richstone (1995) first found a correlation between $M_\bullet$ and the mass of the spheroidal component, M_{bul} . In a comprehensive study using ground-based spectroscopy and HST imaging, Magorrian *et al.* (1998), presented strong evidence in support of the $M_\bullet - M_{bul}$ relation. More recently, two studies (Gebhardt *et al.* 2000a; Ferrarese & Merritt 2000) reported relationships between $M_\bullet$ and σ_e , the stellar velocity dispersion obtained within a large aperture extending to the galaxy effective radius, thus insensitive to the influence of the black hole. The correlation between these two parameters is remarkably strong suggesting a link between the formation of the bulge and the development of the black hole.

In this paper we combine reverberation mapping measurements of $M_\bullet$ in AGN with Narrow Line Region (NLR) gas and bulge stellar kinematic measurements corresponding to motion well beyond influence of the black hole to compare the nuclei of active and normal galaxies. We also investigate the possibility of a correlation between black hole mass and the radio luminosity in AGN. Section 2 describes the data collected from the literature. Section 3 presents the $M_\bullet - \sigma_*$ relation for AGN. In section 4 we

consider the relationship between $M_\bullet$ and the radio luminosity and in Section 5 we summarize our conclusions.

2. DATA

We start with all AGN with black hole masses, measured by the reverberation mapping technique: 17 Seyfert 1 galaxies (as compiled by Wandel *et al.* 1999) and 17 PG quasars (Kaspi *et al.* 2000). For most of the galaxies, we use the black hole masses tabulated by Kaspi *et al.* (2000) evaluated using the mean $H\beta$ profile width. $M_\bullet$ values for three galaxies (Mkn 279, NGC 3516, and NGC 4593) were taken from Ho (1999). For as many objects as possible we obtained published measurements of σ_* , $\text{FWHM}_{[\text{OIII}]}$, and the radio flux. The adopted values are presented in Table 3. For the Seyferts, $\text{FWHM}_{[\text{OIII}]}$ is taken from Whittle (1992a). For the quasars $\text{FWHM}_{[\text{OIII}]}$ values were only adopted if the spectroscopy was of medium to high resolution ($R > 1500$). σ_* in AGN host galaxies has been determined using spectral regions around both the Ca II triplet and Mg*b*. The values used here have been published in Smith, Heckman & Illingworth (1990), Terlevich, Diaz & Terlevich (1990) and Nelson & Whittle (1995) (see Nelson & Whittle 1995, for references on individual galaxies). Radio luminosities are from Whittle (1992a) for Seyferts and from Kellerman *et al.* (1989) for PG quasars. We have scaled observations at 5GHz assuming a power-law index of 0.7 and assume $H_0 = 80\text{kms}^{-1}\text{Mpc}^{-1}$. We also calculate R , the ratio of the radio to optical flux, using the mean continuum luminosities from Kaspi *et al.* (2000).

We point out that the stellar and gas kinematic measurements are from small aperture observations typically covering $2 - 3''$, somewhat smaller than the apertures used to measure σ_e . Nevertheless, we can be confident that these measurements are unaffected by the presence of the black hole. First, since the AGN are at larger distances than the normal galaxies for which $M_\bullet$ has been determined, the aperture typically corresponds to physical scales of a kpc or more, actually comparable to the bulge effective radius. Second Nelson & Whittle (1996) found that Seyferts were offset from the Faber-Jackson relation ($M_{\text{bul}} \propto \log \sigma_*$) for normal galaxies in the sense of having lower velocity dispersions than normal galaxies of the same M_{bul} . Their interpretation was that Seyfert bulges have lower M/L ratios than normal spirals, exactly the opposite of what one would expect if the stellar kinematics were strongly influenced by a massive nuclear black hole.

3. THE $M_\bullet - \sigma_*$ RELATION FOR AGN

Recently, Gebhardt *et al.* (2000b) included Seyferts in their plot of σ_* vs. $M_\bullet$ (in some cases choosing different values for σ_* than in Table 3) and conclude that AGN show no significant difference from the overall relation. Unfortunately, σ_* values are available in the literature for only a hand-full of the AGN with reverberation mapping estimates for $M_\bullet$. Therefore a definitive comparison of AGN and normal galaxies is not possible with these data. However, Nelson & Whittle (1996) have shown that for the majority of Seyfert galaxies, a moderately strong correlation between $\text{FWHM}_{[\text{OIII}]}$ and σ_* exists indicating roughly equal absorption and emission line widths. Thus the $[\text{OIII}] \lambda 5007$ profiles are dominated by virial motion in the bulge potential. As might be expected, a fair amount of real (*i.e.*

not due to measurement error) scatter does exist. Nelson & Whittle (1996) used the deviation from purely virial gas motion to investigate possible secondary influences on the NLR kinematics. Their results confirmed conclusions from previous studies (*e.g.* Whittle 1992b,c) that the interaction of NLR gas with a relatively strong kiloparsec-scale linear radio source can produce non-virial gas acceleration. They also found a weak tendency for interacting systems and mergers to have broader emission lines than expected from purely virial motion, an issue that may be important in quasars which are more likely to be interacting systems (McLeod & Rieke 1995). Thus, keeping the issue of secondary influences on the NLR kinematics in mind, we can proceed with the idea that the primary influence on the forbidden line widths in AGN is the bulge potential.

Table 1
AGN with Measured Black Hole Masses

Name	$\text{Log } M_\bullet$ $M_\odot$	σ_* km s^{-1}	$[\text{OIII}]$ FWHM km s^{-1}	$\text{Log } L_{1415}$ W Hz^{-1}	R	$[\text{OIII}]$ Ref.
3C 120	7.36	162	230	23.72	16.8	8
3C 390.3	8.53	—	410	24.74	198.2	4
AKN 120	8.26	—	490	22.16	0.25	8
F 9	7.90	—	425	—	—	8
IC 4329A	6.70	—	550	22.85	10.2	8
MKN 79	7.72	—	350	22.13	0.8	8
MKN 110	6.75	—	290	22.40	1.6	8
MKN 279	7.62	—	580	22.56	—	8
MKN 335	6.80	—	280	21.68	0.2	8
MKN 509	7.76	—	520	22.43	0.4	8
MKN 590	7.25	169	400	22.05	0.5	8
MKN 817	7.64	—	330	22.28	0.8	8
NGC 3227	7.59	144	485	21.60	4.7	6
NGC 3516	7.36	235	250	21.54	—	6
NGC 3783	6.97	—	230	21.66	0.6	8
NGC 4051	6.11	88	190	21.12	6.0	6
NGC 4151	7.18	171	425	22.23	5.6	6
NGC 4593	6.91	124	255	20.50	—	6
NGC 5548	8.09	—	410	22.22	1.5	6
NGC 7469	6.81	—	360	22.85	3.0	8
PG 0026+129	7.73	—	300	23.71	1.6	7,3
PG 0052+251	8.34	—	—	22.95	0.3	—
PG 0804+761	8.28	—	800	22.64	0.2	—
PG 0844+349	7.33	—	534	21.79	0.1	—
PG 0953+414	8.26	—	640	23.77	1.1	—
PG 1211+143	7.61	—	410	22.47	0.1	5
PG 1226+023	8.74	—	—	27.67	1621.2	—
PG 1229+204	7.88	—	260	22.12	0.3	4
PG 1307+085	8.45	—	458	22.63	0.2	1
PG 1351+640	7.66	—	540	22.29	—	3
PG 1411+442	7.90	—	635	22.38	0.2	—
PG 1426+015	8.67	—	—	22.61	0.2	—
PG 1613+658	8.38	—	585	23.41	0.8	4
PG 1617+175	8.44	—	—	22.85	0.7	—
PG 1700+518	7.78	—	—	25.05	8.5	—
PG 1704+608	7.57	—	425	27.00	562.8	2
PG 2130+099	8.16	—	349	22.56	0.4	7

$[\text{OIII}]$ FWHM References: 1 Appenzeller & Wagner (1991), 2 Brotherton *et al.* (1996), 3 Gelderman & Whittle (1996), 4 Heckman, Miley & Green (1984), 5 Marziani *et al.* (1996), 6 NW95, 7 Vrtilik & Carleton (1985) (1985), 8 Whittle (1992a)

In Figure 1 we plot $\text{FWHM}_{[\text{OIII}]} / 2.35$ or σ_* vs. $M_\bullet$. Filled symbols show the $\text{FWHM}_{[\text{OIII}]}$ values for the galaxies in Table 3, $\bullet$ for Seyferts, $\blacksquare$ for the PG quasars, + signs show AGN for which σ_* measurements exist and triangles show $M_\bullet$ and σ_e from Gebhardt *et al.* (2000a). The similarity in the trends between the data from Table 1 and the Gebhardt *et al.* (2000a) sample is striking. The long solid line is the relation determined by Gebhardt *et al.* (2000a) and the shorter dashed line is our fit to

the AGN, using the ordinary least-squares bisector (Isobe *et al.* 1990) which gives good results for large uncertainties in both variables. The correlation for the AGN is moderately strong ($R = 0.51$, $P(\text{null}) = 0.5\%$) with a slope 3.7 ± 0.7 and intercept -0.5 ± 0.1 . The AGN relation is slightly shifted to larger widths for a given $M_\bullet$ than the Gebhardt *et al.* (2000a) relation and is more scattered. Also, there seems to be no statistical difference between the distributions of Seyferts and quasars.




FIG. 1.— The data from Gebhardt *et al.* (2000a) is shown as $\triangle$ and the solid line is the $M_\bullet - \sigma_e$ relation derived there. Objects with σ_* from Nelson & Whittle (1996) are shown as $+$. Objects for which the [OIII] line widths have been converted to σ are shown as $\bullet$ for Seyferts and as $\blacksquare$ and the dashed line is the fit.

The slope on the fit agrees well with that found by Gebhardt *et al.* (2000a) who derived a different value than Ferrarese & Merritt (2000) (3.75 as opposed to 4.8). Gebhardt *et al.* (2000a) discuss various explanations for the difference in the two values. Although our result supports a shallower slope, an analysis using stellar velocity dispersion instead of emission line width would be more conclusive.

We can be more quantitative about the similarity of the relation for elliptical galaxies and that for the AGN, by considering the origin of the scatter. The mean $1 - \sigma$ error bar in $\log M_\bullet$, plotted in the upper left, is $+0.17 / -0.20$ while the r.m.s. deviation of the filled symbols in Figure 1, relative to the Gebhardt *et al.* (2000a) relation is 0.55. Since we expect that emission line widths can be determined to accuracies of $\sim 10\%$ for the spectral resolution of the observations, we expect that a large component of the scatter is not due to measurement error but reflects real deviations perhaps indicating secondary influences on NLR kinematics. If we calculate the difference from the Gebhardt *et al.* (2000a) relation, this time in $\log \text{FWHM}_{[\text{OIII}]}$, we find a small offset, $\Delta \log \text{FWHM}_{[\text{OIII}]} = 0.06$, with a distribution of r.m.s. width 0.14. Nelson & Whittle (1996) used differences in emission and absorption line widths, $\Delta W \equiv \log \text{FWHM}_{[\text{OIII}]} - \log 2.35 \sigma_*$, to investigate non-virial influences on emission line kinematics. For their entire Seyfert sample they found $\Delta W = 0.0$ and an r.m.s. width of 0.2. Excluding interacting galaxies and objects with relatively luminous linear radio sources they found

$\Delta W = -0.1$, thus slightly narrower emission lines than absorption lines, and an r.m.s. width of 0.13. Thus it is reasonable that the scatter and the small offset indicate non-virial contributions to the line widths in some objects. The fact that the r.m.s. deviation is comparable to that found in Nelson & Whittle (1996) is consistent with this interpretation. Unfortunately, the current sample includes no Seyferts with strong linear radio sources and, due to their larger distances, the radio maps for the quasars are mostly unresolved on kpc scales (see *e.g.* Kellerman *et al.* 1994; Kukula 1998). Therefore, no strong statements regarding non-virial gas kinematics can be made.

The relationship between $M_\bullet$ and the $\text{FWHM}_{[\text{OIII}]}$ in AGN is precisely what we would expect if AGN follow the $M_\bullet - \sigma_*$ relation and if the primary influence on NLR kinematics is the bulge gravitational potential. Although one might point out that the AGN plotted with σ_* do seem to lie systematically below the relation, the reality of such a shift based on only 7 objects is not compelling. Our primary conclusion is that the relationship between the mass of the central dark object and the depth of the gravitational potential in AGN is the same as for normal galaxies. Clearly more velocity dispersion measurements for AGN are needed. However, in fainter objects detailed studies of the [OIII] emission line profiles accounting for non-virial kinematic components could provide a valuable substitute.




FIG. 2.— $\log L_{1415 \text{ MHz}}$ vs. $M_\bullet$ for AGN.

These results seem to conflict with previous reports that AGN have lower black hole to bulge mass ratios than normal galaxies (Wandel *et al.* 1999) or alternatively that the reverberation mapping estimates of $M_\bullet$ are systematically low (Ho 1999). Figure 1 suggests that neither of these is correct and that the differences must lie in the values of M_{bul} . A number of factors may have contributed to this apparent disagreement. First, the $M_\bullet - \sigma_e$ relation derived by Gebhardt *et al.* (2000a) has been evaluated using only stellar dynamical black hole mass estimates from three-integral modeling. The result is that the black hole masses are about a factor of 3 lower than the ones published by Magorrian *et al.* (1998) which were used in Wandel *et al.* (1999). Second Nelson & Whittle (1996) found that Seyferts are offset from the Faber-Jackson relation

for normal galaxies having brighter bulges by on average about 0^m.6 at the same velocity dispersion. More generally bulge-disk decomposition is a difficult task even for normal, nearby galaxies. The problem is compounded in Seyfert 1 galaxies by their smaller apparent bulge sizes, due to larger mean distances, and the bright point source component. Thus, much of the discrepancy can be resolved by acknowledging that as much or greater uncertainty exists in the bulge luminosity of Seyferts as exists in the estimates of $M_{\bullet}$ by any of the available techniques.

4. RADIO LUMINOSITY AND BLACK HOLE MASS IN AGN

Using reliable measurements of black hole mass in AGN we can compare parameters related to the energetics of AGN with the size of the central engine. For example Kaspi *et al.* (2000) found a strong correlation between the black hole mass and the mean optical continuum luminosity, with the dependence $M \propto L^{0.5}$, suggesting that more luminous AGN radiate at a larger fraction of the Eddington luminosity. Nelson & Whittle (1996) found a correlation between σ_* and the radio luminosity at 1415 MHz, L_{1415} , in Seyfert galaxies. This relation was linked to previously known correlations between radio power and absolute blue luminosity (Meurs & Wilson 1984; Edelson 1987; Whittle 1992c) and between radio power and [OIII] line width (*e.g.* Whittle 1985). They speculated that the origin of such a correlation might result from the correlation of $M_{\bullet}$ with bulge mass and that the radio sources associated with larger central engines might be correspondingly more luminous. In Figure 2 we plot $L_{1415\text{MHz}}$ vs. $M_{\bullet}$, plotting different symbols for radio-loud ($R > 8.0, +$), radio-intermediate ($1.0 < R < 8.0, \circ$) and radio-quiet AGN ($R < 1.0, \bullet$). The criteria are cho-

sen rather arbitrarily, but nevertheless demonstrate some interesting differences. The solid line shows a fit to the most radio-quiet group. The correlation is quite strong ($R = 0.82$, $P(\text{null})=0.003\%$) with a slope 0.84 ± 0.12 . A fit including radio-intermediate data (not shown) is somewhat weaker ($R = 0.58$, $P(\text{null})=0.1\%$) and a fit to the entire sample weaker still ($R = 0.40$, $P(\text{null})=2\%$). The results suggest that at least in radio-quiet AGN the radio luminosity, emitted on kpc scales is linked to the mass of the nuclear black hole. Nelson & Whittle (1996) noted other explanations for their result, including higher ambient pressure in larger bulges leading to increased radio emissivity. We cannot distinguish between these possibilities and it is possible that both effects work together to produce the observed correlation.

5. SUMMARY

Using the assumption that NLR kinematics are predominantly due to virial motion in the host galaxy potential, we find that AGN follow a relation between $M_{\bullet}$ and σ_e similar to normal galaxies. Although the AGN black hole masses are determined using reverberation mapping, we find no significant differences with results from studies using stellar dynamical techniques. Satisfied that the black hole masses in AGN are reliable we can begin to examine the relationship of black hole mass with other properties of AGN. As an example, we have found that the radio luminosity in radio-quiet AGN is correlated with black hole mass relating parameters determined on vastly different size scales.

I would like to thank Donna Weistrop for her support and comments on a draft of this letter. This work was supported in part by NASA, under contract NAS5-31231.

REFERENCES

- Appenzeller, I., & Wagner, S. J. 1991 A&A, 250, 57
 Bower, G. A., Green, R. F., Danks, A., Gull, T., Heap, S., Hutchings, J., Joseph, C., Kaiser, M. E., Kimble, R., Kraemer, S., Weistrop, D., Woodgate, B., Lindler, D., Hill, R. S., Malumuth, E. M., Baum, S., Sarajedini, V., Heckman, T. M., Wilson, A. S. & Richstone, D. O. 1998, ApJ, 522, L111
 Bower, G. A., Green, R. F., Bender, R., Gebhardt, K., Lauer, T. R., Magorrian, J., Richstone, D. O., Danks, A., Gull, T., Hutchings, J., Joseph, C., Kaiser, M. E., Weistrop, D., Woodgate, B., Nelson, C., Malumuth, E., & Valluri, M. 2000, ApJ, in press.
 Brotherton, M. S. 1996, ApJS, 102, 1
 Edelson, R. A. 1987, ApJ, 313, 651
 Ferrarese, L. & Merritt, D. 2000, ApJ, 539, L9
 Gebhardt, K., Bender, R., Bower, G., Dressler, A., Faber, S. M., Filippenko, A. V., Green, R., Grillmair, R., Ho, L. C., Kormendy, J., Lauer, T. R., Magorrian, J., Pinkney, J., Richstone, D., Tremaine, S. 2000, ApJ, 539, 13
 Gebhardt, K., Bender, R., Bower, G., Dressler, A., Faber, S. M., Filippenko, A. V., Green, R., Grillmair, R., Ho, L. C., Kormendy, J., Lauer, T. R., Magorrian, J., Pinkney, J., Richstone, D., Tremaine, S. 2000, ApJ, in press
 Harms, R. J., Ford, H. C., Tsvetanov, Z. I., Hartig, G. F., Dressler, L. L., Kriss, G. A., Bohlin, R., Davidsen, A. F., Margon, B., Kochhar, Ajay K. 1994, ApJ, 435, 35
 Heckman, T. M., Miley, G. K., & Green, R. F. 1984, ApJ, 281, 525
 Ho, L. C. 1999, in *Observational Evidence for Black Holes in the Universe*, ed. S. K. Chakrabarti (Dordrecht:Kluwer), 153
 Gelderman, R. F., & Whittle, M. 1994, ApJS, 91, 491
 Isobe, T., Feigelson, E. D., Akritas, M. G., & Babu, G. J., 1990, ApJ, 364, 104
 Kaspi, S., Smith, P. S., Netzer, H., Maoz, D., Januzzi, B. T., & Givon, U. 2000, ApJ, 533, 631
 Kellerman, K. I., Sramek, R. A., Schmidt, M., Shaffer, D. B., & Green, R. 1989, AJ, 98, 1195
 Kellerman, K. I., Sramek, R. A., Schmidt, M., Green, R. F., & Shaffer, D. B. 1994, AJ, 108, 1163
 Kormendy, J. & Richstone, D. 1995, ARA&A, 33, 581
 Kormendy, J. 1988, ApJ, 335, 40
 Kukula, M. J., Dunlop, J. S., Hughes, D. H., & Rawlings, S. 1998, MNRAS, 297, 366
 Magorrian, J., Tremaine, S., Richstone, D., Bender, R., Bower, G., Dressler, A., Faber, S. M., Gebhardt, K., Green, R., Grillmair, C., Kormendy, J., & Lauer, T. 1998, AJ, 115, 2285
 Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., & Moles, M. 1996, ApJS, 104, 37
 McLeod, K. K., & Reike, G. 1994, ApJ, 431, 137
 Meurs, E. J. A., & Wilson, A. S. 1984, A&A, 136, 206
 Nelson, C. & Whittle, M. 1995, ApJS, 99, 67
 Nelson, C. & Whittle, M. 1996, ApJ, 465, 96
 Netzer, H., & Peterson, B. M. 1997, in *Astronomical Time Series*, ed. D. Maoz, A. Sternberg, & E. M. Leibowitz (Dordrecht: Kluwer), 85
 Peterson, B. & Wandel, A. 1999, ApJ, 521, 95
 Smith, E. P., Heckman, T. M. & Illingworth, G. D. 1990, ApJ, 356, 399
 Terlevich, E., Diaz, A. I. & Terlevich, R. 1990, MNRAS, 242, 271
 Vrtillek, J. M., & Carleton, N. P. 1985, ApJ, 294, 106
 Wandel, A., Peterson, B. M., & Malkan, M. A. 1999, ApJ, 519, L39
 Wandel, A. 1999, ApJ, 526, 579
 Whittle, M. 1985, MNRAS, 213, 33
 Whittle, M. 1992, ApJS, 79, 49
 Whittle, M. 1992, ApJ, 387, 109
 Whittle, M. 1992, ApJ, 387, 121